

REMARKS ON NON-COMPACT COMPLETE RICCI EXPANDING SOLITONS

LI MA AND DEZHONG CHEN

ABSTRACT. In this paper, we study gradient Ricci expanding solitons (X, g) satisfying

$$Rc = cg + D^2f,$$

where Rc is the Ricci curvature, $c < 0$ is a constant, and D^2f is the Hessian of the potential function f on X . We show that for a gradient expanding soliton (X, g) with non-negative Ricci curvature, the scalar curvature R has at least one maximum point on X , which is the only minimum point of the potential function f . Furthermore, $R > 0$ on X unless (X, g) is Ricci flat. We also show that there is exponentially decay for scalar curvature for ϵ -pinched complete non-compact expanding solitons.

1. INTRODUCTION

In this paper, we continue our study on Ricci solitons [7], which are generated by one parameter family of diffeomorphisms and are special solutions to Ricci flow introduced by R. Hamilton in 1982 [?]. We assume in this paper that (X, g) is a gradient expanding soliton. Let recall the definition of expanding soliton.

Definition 1. *We call a Riemannian manifold (X, g) an expanding soliton if there is a smooth solution f on a Riemannian manifold (X, g) such that for some constant $c < 0$, it holds the equation*

$$(1) \quad Rc = cg + D^2f,$$

on X , where D^2f is the Hessian matrix of the function f and Rc is the Ricci tensor of the metric g . We call the function f the potential function for the soliton (X, g) . If $c > 0$ in (1), (X, g) is called a shrinking soliton; if $c = 0$, (X, g) is called a steady soliton.

In the study of Ricci flow, we often meet the following definition.

Date: July. 20th, 2005.

1991 Mathematics Subject Classification. 53Cxx.

Key words and phrases. Ricci flow, expanding soliton.

* This work is supported in part by the Key 973 project of Ministry of Science and Technology of China.

Definition 2. *The Ricci curvature of a Riemannian manifold (X, g) is called ϵ -pinched if there is some $\epsilon > 0$ such that the scalar curvature $R > 0$ on X and*

$$Rc \geq \epsilon Rg$$

on X .

Throughout this paper, we shall assume that the Riemannian manifold (X, g) is a complete non-compact Riemannian manifold of dimension $n \geq 3$. We denote by R the scalar curvature of the metric g .

Our main result is the following

Main Theorem . *Assume that the Ricci curvature of the gradient expanding soliton (X, g) is non-negative. Then the scalar curvature R has at least one maximum point on X , which is the only minimum point of the potential function f . Furthermore, $R > 0$ on X unless (X, g) is Ricci flat.*

The proof of this Theorem will be proved in section 3.

In section four, we will prove the following result

Theorem 3. *Assume that (X, g) is a gradient expanding soliton with its Ricci curvature being ϵ -pinched. Then its scalar curvature has the decay*

$$R(s) \leq R(o)e^{Cs-Cs^2}.$$

as the distance function s from a fixed point going to infinity, i.e., $s = d(x, o) \rightarrow +\infty$.

We remark that a similar but weaker decay result is announced by L.Ni in Proposition 3.1 in [8]. We know the result for a while and a reason for the delay of this present is that we try to prove non-existence of this kind of expanding solitons. However, we have not been succeed yet.

Throughout C will denote various uniform constants in different places.

2. PRELIMINARY

We recall first some basic properties about Ricci solitons [6].

Taking the trace of both sides of (1), we have

$$(2) \quad R = nc + \Delta f.$$

Take a point $x \in X$. In local normal coordinates (x^i) of the Riemannian manifold (X, g) at a point x , we write the metric g as (g_{ij}) .

The corresponding Riemannian curvature tensor and Ricci tensor are denoted by $Rm = (R_{ijkl})$ and $Rc = (R_{ij})$ respectively. Hence,

$$R_{ij} = g^{kl} R_{ikjl}$$

and

$$R = g^{ij} R_{ij}.$$

We write the covariant derivative of a smooth function f by $Df = (f_i)$, and denote the Hessian matrix of the function f by $D^2f = (f_{ij})$, where D the covariant derivative of g on X . The higher order covariant derivatives are denoted by f_{ijk} , etc. Similarly, we use the $T_{ij,k}$ to denote the covariant derivative of the tensor (T_{ij}) . We write $T_j^i = g^{ik} T_{jk}$. Then the Ricci soliton equation is

$$R_{ij} = f_{ij} + cg_{ij}.$$

Taking covariant derivative, we get

$$f_{ijk} = R_{ij,k}.$$

So we have

$$f_{ijk} - f_{ikj} = R_{ij,k} - R_{ik,j}.$$

By the Ricci formula we have that

$$f_{ijk} - f_{ikj} = R_{ijk}^l f_l.$$

Hence we obtain that

$$R_{ij,k} - R_{ik,j} = R_{ijk}^l f_l.$$

Recall that the contracted Bianchi identity is

$$R_{ij,j} = \frac{1}{2} R_i.$$

Upon taking the trace of the previous equation we get that

$$\frac{1}{2} R_i + R_i^k f_k = 0,$$

i.e.,

$$(3) \quad R_k = -2R_k^j f_j.$$

Then at x ,

$$D_k(|Df|^2 + R + 2cf) = 2f_j(f_{jk} - R_{jk} + 2cg_{jk}) = 0.$$

So,

$$(4) \quad |Df|^2 + R + 2cf = M,$$

where M is a constant.

In the remaining part of this section, we assume that $0 \leq Rc \leq C$ on the expanding soliton (X, g) for some constant $C > 0$. Then we have

$|D^2 f| \leq C$ on X . Assume $f \geq 0$ and that o is a critical point of the potential function f . Then using the Taylor's expansion, we have

$$f(x) \leq C d^2(x, o).$$

We now study the behavior of the potential function along a minimizing geodesic curve on the expanding soliton. A similar work has been done by G. Perelman [9] (see also [5]) where he tries to give some uniform bounds on potential function f on a shrinking soliton. Fix a point $o \in X$. Take any minimizing geodesic curve $\gamma(s)$ connecting x and the fixed point p , where s is the arc-length parameter. Write by $r = d(x, o)$ and $X = \gamma'(s)$. Assume that $r > 2$. Let $\{Y_i\}$ ($i = 1, \dots, n-1$) be an orthonormal parallel vector fields along γ . Let Y be an orthogonal vector field along the curve γ vanishing at end points. Then the second variational formula [10] (see also [1]) tells us that

$$\int_0^r (|Y|^2 - \langle R(X, Y)Y, X \rangle) ds \geq 0.$$

Take Y to be sY_i on $[0, 1]$, $= Y_i$ on $[1, r - r_0]$ where $1 < r_0 < r$, and $\frac{r-s}{r_0}Y_i$. Adding over i gives that

$$\int_0^r Rc(X, X) \leq C_0(r_0) + \frac{n-1}{r_0} - \int_{r-r_0}^r \left(\frac{r-r_0}{r_0}\right)^2 Rc(X, X) ds,$$

which implies that for some constant $C > 0$,

$$(5) \quad \int_0^r Rc(X, X) \leq C.$$

Note that

$$\left(\int_0^r Rc(X, Y_1) ds\right)^2 \leq r \int_0^r |Rc(X, Y_1)|^2 ds \leq s \sum_i \int_0^r |Rc(X, Y_i)|^2 ds.$$

Thinking of Rc as self-adjoint linear operator on TX and taking a point-wise orthonormal frame $\{e_j\}$ as eigenvectors of $Rc = (\bigoplus \lambda_j)$, we have that

$$R = \sum_j \lambda_j$$

and for $X = \sum_j X_j e_j$,

$$\sum_i |Rc(X, Y_i)|^2 = \langle X, Rc^2 X \rangle = \sum_j \lambda_j X_j^2 \leq R Rc(X, X).$$

Then,

$$\left(\int_0^r Rc(X, Y_1) ds\right)^2 \leq C s \int_0^r Rc(X, X) \leq C^2 s.$$

Hence, for any unit vector field Y along γ , orthogonal to X , we have

$$\int_0^r Rc(X, Y) ds \leq C(\sqrt{s} + 1).$$

Using (1) we have

$$\frac{d^2 f(\gamma(s))}{ds^2} = Rc(X, X) - c \geq -c,$$

and

$$\frac{d(Yf)(\gamma(s))}{ds} = Rc(X, Y).$$

Then we have

$$\frac{df(\gamma(s))}{ds} \geq \frac{df(\gamma(s))}{ds}(0) - cs \geq -cs + C$$

and for $s > 2$,

$$(6) \quad |(Yf)(\gamma(s))| \leq |(Yf)(\gamma(0))| + \int_0^s |Rc(X, Y)| ds \leq C\sqrt{s}.$$

Therefore, we can conclude that at large distance from o the potential function f has its gradient making small angle with the gradient of the distance function from o .

3. PROOF OF MAIN THEOREM

Assume that $Rc \geq 0$ on X , and we also assume that for some constant $c < 0$ we have $Rc - cg > 0$ on X . By (1) we know that

$$D^2 f = Rc - cg \geq -cg > 0, \quad \text{on } X.$$

Then the potential function f is locally strictly convex. Since (X, g) is a complete non-compact Riemannian manifold, we have that f has at most one critical point, i.e., the point where $\nabla f = 0$. Using $D^2 f > 0$, we know that if $p \in X$ is the critical point of f , then it is a non-degenerate minimum point of f .

Note that along any minimizing geodesic curve $\gamma(s)$ connecting x and the fixed point p , where s is the arc-length parameter, we have

$$(7) \quad \begin{aligned} \langle \nabla f, \gamma'(s) \rangle|_0^s &= \int_0^s f_{ij} \frac{d\gamma^i}{ds} \frac{d\gamma^j}{ds} ds \\ &= \int_0^s (R_{ij} - cg_{ij}) \frac{d\gamma^i}{ds} \frac{d\gamma^j}{ds} ds \\ &= -cs + \int_0^s R_{ij} \frac{d\gamma^i}{ds} \frac{d\gamma^j}{ds} ds \\ &\geq -cs > 0 \end{aligned}$$

This implies that $f(\gamma(s))$ is growing at infinity at least the quadratic rate $-c$ of the distance function. Then f has at least a minimum point in X .

Assume that o is the only critical point of f . Then by adding a constant, we can assume that $f(o) = 0$ and $f > 0$ on $X - \{o\}$. Using (4), we know that

$$M = |Df|^2(o) + R(o) + 2cf(o) = R(o).$$

Using (3) we know that o is also the critical point of R .

Let $x \in X - \{o\}$. Taking a minimizing geodesic curve $\gamma(s)$ connecting x and the fixed point o , where s is the arc-length parameter, we again have by using (7)

$$\langle \nabla f, \gamma'(s) \rangle > -cs > 0.$$

This implies that the integral curves of ∇f in $X - \{o\}$ emanating from the point o to infinity. Take a integral curve $\sigma(t)$ ∇f in $X - \{o\}$. Then by (3) we have

$$(8) \quad \frac{d}{dt}R(\sigma(t)) = R_i f_i = -2Rc(\nabla f, \nabla f) \leq 0.$$

Hence $R(x) \leq R(o)$ for all $x \in X - \{o\}$. So, o is a maximum point of R .

By this we conclude that

Assertion 4. *Assume that the Ricci curvature of the gradient expanding soliton (X, g) is non-negative positive. Then the scalar curvature R has at least one maximum point of R , which is the only critical point of the potential function f .*

If $R(o) = 0$, then $R = 0$ on X . Hence $Rc = 0$ on X , that is to say that (X, g) is Ricci flat. So we have $R(o) > 0$. By the local strong maximum principle, we must have $R > 0$ on the whole space X .

This finishes the *proof of Main Theorem*.

In the remaining part of this section, we consider the behavior of f at infinity. Since

$$|Df|(x)^2 + 2cf(x) = R(o) - R(x) \geq 0,$$

we get that

$$|Df|^2 \geq -2cf = 2|c|f.$$

Then we have

$$|D\sqrt{f}| \geq \sqrt{\frac{|c|}{2}},$$

at where $f \neq 0$. Therefore, we have

$$\sqrt{f}(s) \geq \sqrt{\frac{|c|}{2}}s$$

and

$$f(s) \geq \frac{|c|}{2}s^2$$

along any minimizing geodesic curve $\gamma(s)$ connecting x and the fixed point o , where s is the arc-length parameter.

Note that using (2) we have

$$|Df|^2(s) = -2cf(x) + R(o) - R(x) \leq -2cf(x) + R(o) \leq Cs^2 + R(o).$$

Hence, for $s \gg 1$,

$$(9) \quad C_4s \leq |Df|(s) \leq C_5s.$$

4. ϵ PINCHED SOLITONS

We give a proof of Theorem 3 below. We try to make the proof more transparent and self-contained.

Proof of Theorem 3: Recall that the Ricci curvature of the non-shrinking soliton (X, g) is ϵ -pinched, i.e., for some $\epsilon > 0$ we have that $R > 0$ on X and

$$Rc \geq \epsilon Rg$$

on X . Then using the maximum principle, we know that either $R = 0$ on X or $R > 0$. If $R = 0$ on X , then by the pinching condition we know that (X, g) is Ricci flat.

Assume that $R > 0$ on X . Then as before, the potential function f is locally strictly convex. Since (X, g) is a complete non-compact Riemannian manifold, we have that f has at most one critical point, i.e., the point where $\nabla f = 0$. Assume that we have a critical point for f , saying that it is $o \in X$. Then using (3), we know it is also a critical point of R . Using (8), we know that is the maximum point for R . In particular, we know that R is a bounded function on X , saying that $D > 0$ is the upper bound.

Using (3) and the ϵ -pinched condition, we have that

$$-R|\nabla f|^2 \leq \langle \nabla R, \nabla f \rangle = -2Rc(\nabla f, \nabla f) \leq -\epsilon R|\nabla f|^2.$$

Taking a minimizing geodesic curve $\gamma(s)$ connecting x and a fixed point o , where s is the arc-length parameter, we have

$$\begin{aligned} \langle \nabla f, \gamma'(s) \rangle \Big|_0^s &= \int_0^s f_{ij} \frac{d\gamma^i}{ds} \frac{d\gamma^j}{ds} ds \\ &= \int_0^s (R_{ij} - cg_{ij}) \frac{d\gamma^i}{ds} \frac{d\gamma^j}{ds} ds \\ (10) \quad &= -cs + \int_0^s R_{ij} \frac{d\gamma^i}{ds} \frac{d\gamma^j}{ds} ds. \end{aligned}$$

This implies that there is a constant C_2 such that

$$\langle \nabla f, \gamma'(s) \rangle \geq -cs + \int_0^s \epsilon R ds \geq -cs + \int_0^1 R ds \geq -cs + C_2 \geq C_2$$

for $s \gg 1$.

Using (5) and the pinching condition, we have that

$$\int_0^s R ds \leq C_6.$$

Using the pinching condition again, (10) also implies that

$$\langle \nabla f, \gamma'(s) \rangle \leq -cs + \int_0^s R ds \leq -cs + D.$$

Therefore, the angle between ∇f and the gradient of the distance function from o is almost fixed.

Then, using (3) and the ϵ -pinched condition, we have for some constant $C_3 > 0$,

$$(R^{-1})_s = -R^{-2} \langle \nabla R, \gamma'(s) \rangle = 2R^{-2} Rc(\nabla f, \gamma'(s)).$$

Using (6) and (9), we obtain that

$$\begin{aligned} Rc(\nabla f, \gamma'(s)) &= |\nabla f| Rc(\gamma', \gamma') + 0(\sqrt{s}) \\ &\geq \epsilon R |\nabla f| + 0(\sqrt{s}) \geq R(Cs - C), \end{aligned}$$

we have

$$(R^{-1})_s \geq 2R^{-1}(Cs - C).$$

This implies that

$$(\log R)_s \leq C - Cs$$

and

$$R(s) \leq R(o)e^{Cs - Cs^2}.$$

This implies that $R \rightarrow 0$ exponentially as $s \rightarrow +\infty$. *This completes the proof of Theorem 3.*

Theorem 3 tells us that for such (X, g) we have

$$A = \limsup_{s \rightarrow \infty} Rs^2 = 0.$$

REFERENCES

- [1] Th. Aubin, *Non-linear Analysis on manifolds*, Springer, New York, 1982.
- [2] S. Bando, A. Kasue, and H. Nakajima, *On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth*, Invent. Math., 97(1989)313-349.
- [3] R. Bryant, *Gradient Kahler-Ricci solitons*, ArXiv.math.DG/0407453, 2004.
- [4] H. D. Cao, *Existence of gradient Kahler-Ricci soliton*, Elliptic and parabolic methods in geometry, Eds. B. Chow, R. Gulliver, S. Levy, J. Sullivan, A K Peters, pp.1-6, 1996.

- [5] B.Kleiner, and J.Lott, *Notes on Perelman's papers*,
<http://www.math.lsa.umich.edu/research/ricciflow/perelman.html>
- [6] R.Hamilton, *The formation of Singularities in the Ricci flow*, Surveys in Diff. Geom., Vol.2, pp7-136, 1995.
- [7] Li Ma, *Remarks on Ricci solitons*, Arxiv.math.DG/0411426, 2004.
- [8] L.Ni, *Ancient solutions to Kahler-Ricci flow*, Math. Research letters. 11(2005)10001-10020.
- [9] G.Perelman, *Ricci flow with surgery on three manifolds*, arXiv:math.Dg/0303109.
- [10] R.Schoen and S.T.Yau, *Lectures on Differential Geometry*, IP, Boston, 1994.
- [11] S.T.Yau, *Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry*, Indiana University Math. Journal, 25(1976)659-670.

DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING
100084, CHINA

E-mail address: lma@math.tsinghua.edu.cn